


FytoSave is for the control of powdery mildew on a range of fruits and vegetables



FytoSave contains 12.5 g/l COS-OGA

Keep out of reach of children.

Do not eat, drink or smoke when using this product.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

To avoid risk to human health and the environment, comply with the instructions for use.

MAPP 18433

Gowan Crop Protection Ltd, Highlands House, Basingstoke Road, Spencers Wood, Reading, Berks, RG7 1NT. (Registered company number: 9852199)

Tel:+44 (0) 1582 280390

24 HOUR EMERGENCY TELEPHONE NUMBER:

+44 (0) 870 820 04 18



Authorisation holder: FytoFend S.A.

FytoSave® is a registered Trademark of FytoFend S.A.

Shake well before use

Lot No.: See Can

Pack Size: 5 L e

74626-UK-FYTF-LF-04-0519

IMPORTANT INFORMATION

FOR USE ONLY AS A PROFESSIONAL FUNGICIDE, AS DIRECTED BELOW

Crops	Protected or Field	Max. individual dose	Maximum number of treatments: (per crop)	Time of application
Aubergine, tomato, pepper and chilli (permanent protection with full enclosure)	Р	2 l/ha	5	3 fully expanded leaves – complete ripening: fruits reach their typical full maturity colour (BBCH 13-89)
Courgette, summer squash, cucumber, melon, winter squash and pumpkin (permanent protection with full enclosure)	Р	2 l/ha	5	3 fully expanded leaves - third fruit has reached its size and typical shape (BBCH 13 - 73)

Other Specific Restrictions

A minimum interval of 7 days must be observed between applications. Treatment must only be made under 'permanent protection' situations which provide full enclosure (including continuous top and side barriers down to below ground level) and which are present and maintained over a number of years. Reasonable precautions must be taken to prevent access of birds, wild mammals and honey bees to treated crops. To minimise airborne environmental exposure, vents, doors and other openings must be closed during and after application until the applied product has fully settled.

EXTENSION OF USE

These extensions of the authorised use provide for the use of FYTOSAVE in respect of the crops and situations, other tthan those included on the product label (above). No efficacy or phytotoxicity data have been assessed and as such the 'extension of use', is at all times done at the user's choosing, and the commercial risk is entirely theirs.

Crops	Protected or Field	Max. individual dose	Maximum number of treatments: (per crop)	Time of application
Table and wine grape	F	2 l/ha	8	-
Strawberry	F&P	2.5 l/ha	7	-
Blackberries, raspberries, loganberry and rubis hybrid	F&P	2.5 l/ha	8	-
Ornamentals	F&P	3 l/ha	8	-
Lettuce, angelica, baby leaf crops, balm, basil, bay, caraway leaves, celery leaves, chervil, chives, coriander, dill leaves, fennel leaves, herbs (others), hyssop, lovage leaves, marjoram, mint, oregano, parsley, rocket, rosemary, sage, salad burnet, savory, sweet cicely, tarragon, thyme	F&P	2 l/ha	8	-
Hops	F	5 l/ha	8	-

Recommended water volumes: Table and wine grapes: 400 to 1000 litres of water per hectare. Ornamental plant production: 1000 litres of water per hectare. Lettuce, rocket, baby leaf crops and various herbs: 500 to 1000 litres of water per hectare. Blackberry, raspberry, loganberry and rubus hybrid and strawberry: 400 to 1000 litres of water per hectare. Hops: 500-1000 litres of water per hectare.

Other Specific Restrictions

This product must only be applied in accordance with the terms of this extension of authorisation, the product label and/or leaflet and any additional guidance on extensions of authorisation. A minimum interval of 7 days between applications is required.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

DIRECTIONS FOR USE

IMPORTANT: This information is authorised as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Pest controlled

Aubergine, tomato, pepper and chilli – moderate control of powdery mildew.

Cucumber, courgette, summer squash, melon, winter squash and pumpkin - control of powdery mildew.

Mixing and spraying

Shake well before use.

Pour Fytosave into the tank half filled with water.

Start agitation during filling and continue until the end of the application. Do not let the mixture remain in the tank.

Water volume:

Aubergine, tomato, pepper and chilli – 750-1000l/ha.

Cucumber, courgette, summer squash, melon, winter squash and pumpkin – 500l/ha.

Further information

The treatment must be carried out preventively before signs of powdery mildew. In the context of the use of FYTOSAVE, powdery mildew management must be done in a preventive way to stimulate the plant before signs of any attack.

It is important to use FytoSave as a preventative treatment and in sequences of 2 to 3 successive applications to obtain good stimulation of the plants defences. The interval between applications should be 7-10 days depending on conditions and disease pressure.

In the case of any established disease or high risk of infection start control with a curative product before applying FYTOSAVE.

Apply FYTOSAVE to healthy crops, avoid application to stressed crops.

SAFETY PRECAUTIONS

Operator protection

Do not re-enter areas/crops before the spray deposit is completely dry.

Environmental protection

DO NOT CONTAMINATE WATER with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

Storage and disposal

- STORE IN A COOL, DRY PLACE in accordance with specific instructions and requirements.
- KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.
- Protect from frost.
- KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.
- EMPTY THE CONTAINER COMPLETELY and dispose of safely.
- DO NOT RE-USE CONTAINER for any purpose.

Medical advices

AFTER INHALATION: Fresh air, rest.

AFTER INGESTION: Rinse mouth, rest.

AFTER SKIN CONTACT: Wash with water and soap, Rinse thoroughly, Remove any contaminated clothing and shoes.

AFTER EYE CONTACT: Rinse with plenty of water for 15 minutes.

In case of symptoms, consult a doctor and show him / her the packaging or the label. Notes for the doctor:

Management: symptomatic treatment.

Approval Holder and Marketing Company: FytoFend S.A. Rue Phocas Lejeune, 25 (B6) 5032 ISNES (Belgium) +32 81 728 840 Container Size: 5 L

BATCH NO.:
Date of manufacture:

74626-UK-FYTF-BL-05-0519